Author Index

Aggarwal, A., Merritt, Jr., J.W. and Nash, T.E.
Cysteine-rich variant surface proteins of *Giardia lamblia*, 39
Aline, Jr., R.F., Scholler, J.K. and Stuart, K.
Transcripts from the co-transposed segment of variant surface glycoprotein genes are in *Trypanosoma brucei* polyribosomes, 169
Anand, R., see Clark, J.T., 15
Anders, R.F., see Favaloro, J.M., 297
Åslund, L., see Franzén, L., 201

Baginsky, W.F., see Schmatz, D.M., 263
Barbet, A.F., Myler, P.J., Williams, R.O. and McGuire, T.C.
Shared surface epitopes among trypanosomes of the same serodeme expressing different variable surface glycoprotein genes, 191
Beach, D.H., see Goad, L.J., 179
Becker, I., see Ostoa-Saloma, P., 133
Berens, R.L., see Goad, L.J., 179
Bisby, R.M., see Browning, P.M., 57
Blanton, R., see Reis, M.G., 113
Brown, K.N., see Viriyakosol, S., 93
Browning, P.M. and Bisby, R.M.
Qinghaosu does not affect the major thermotropic phase transition in model membranes of dipalmitoylphosphatidylcholine, 57
Bülow, R., Nonnengässer, C. and Overath, P.
Release of the variant surface glycoprotein during differentiation of bloodstream to procyclic forms of *Trypanosoma brucei*, 85
Button, L.L., Russell, D.G., Klein, H.L., Medina-Acosta, E., Karess, R.E. and McMaster, W.R.
Genes encoding the major surface glycoprotein in *Leishmania* are tandemly linked at a single chromosomal locus and are constitutively transcribed, 271

Cabrera, N., see Ostoa-Saloma, P., 133
Capron, A., see Dissous, C., 49
Chambers, A.E., see Rodrigues, V., 7
Chaudhri, M., see Rodrigues, V., 7
Clark, J.T., Donachie, S., Anand, R., Wilson, C.F., Heidrich, H.-G. and McBride, J.S.
46–53 Kilodalton glycoprotein from the surface of *Plasmodium falciparum* merozoites, 15
Coggins, J.R., see Jarroll, E.L., 121
Coppel, R.L., see Favaloro, J.M., 297
Coquelet, H., see Paindavoine, P., 61
Crewther, P.E., see Favaloro, J.M., 297

Danforth, H.D., see Jenkins, M.C., 153
Davis, A.H., see Reis, M.G., 113
Delves, C.J., Goman, M., Ridley, R.G., Matile, H., Lensen, T.H.W., Ponnudurai, T. and Scaife, J.G.
Identification of *Plasmodium falciparum*-infected mosquitoes using a probe containing repetitive DNA, 105
Diffley, P. and Mama, K.
Fixed and temporary fluctuations in the cell cycle of monomorphic lines of *Trypanosoma brucei gambiense*, 1
Dissous, C. and Capron, A.
Schistosoma mansoni and its intermediate host *Biomphalaria glabrata* express a common 39 kilodalton acidic protein, 49
Donachie, S., see Clark, J.T., 15

Erlandsen, S.L., see Jarroll, E.L., 121

Favaloro, J.M., Marshall, V.M., Crewther, P.E., Coppel, R.L., Kemp, D.J. and Anders, R.F.
cDNA sequence predicting an octapeptide-repeat antigen of *Plasmodium falciparum* (*Short Communication*), 297
Fetterer, R.H., see Jenkins, M.C., 153
Franzén, L., Wåhlin, B., Wahlgren, M., Åslund, L., Perlmann, P., Wigzell, H. and Pettersson, U.
Enhancement or inhibition of *Plasmodium falciparum* erythrocyte reinvasion in vitro by antibodies to an asparagine rich protein, 201

Garfinkel, L., see Huber, M., 285
Gitler, C., see Huber, M., 285
Goad, L.J., Berens, R.L., Marr, J.J., Beach, D.H. and Holz, Jr., G.G.
The activity of ketoconazole and other azoles against *Trypanosoma cruzi*: biochemistry and chemotherapeutic action in vitro, 179
Goman, M., see Delves, C.J., 105
Gregory, W.F., see Maizels, R.M., 213

Heidrich, H.-G., see Clark, J.T., 15
Holloway, S.P., see Hyde, J.E., 247
Holz, Jr., G.G., see Goad, L.J., 179
Hori, H., see Watanabe, J.-i., 163
Huber, M., Koller, B., Gitler, C., Mirelman, D., Revel, M., Rozenblatt, S. and Garfinkel, L.
Entamoeba histolytica ribosomal RNA genes are carried on palindromic circular DNA molecules, 285
Hundt, E., see Knapp, B., 73
Hyde, J.E., Kelly, S.L., Holloway, S.P., Snewin, V.A. and Sims, P.F.G.
A general approach to isolating *Plasmodium falciparum* genes using non-redundant oligonucleotides inferred from protein sequences of other organisms, 247

Jarroll, E.L., Manning, P., Lindmark, D.G., Coggins, J.R. and Erlandsen, S.L.
Giardia cyst wall-specific carbohydrate: evidence for the presence of galactosamine, 121
Jenkins, M.C., Danforth, H.D., Lillehoj, H.S. and Fetterer, R.H.
cDNA encoding an immunogenic region of a 22 kilodalton surface protein of *Eimeria acervulina* sporozoites, 153
Karess, R.E., see Button, L.L., 271
Kelly, C., see Rodrigues, V., 7
Kelly, C., see Selkirk, M.E., 229

Kelly, S.L., see Hyde, J.E., 247
Kemp, D.J., see Favaloro, J.M., 297
Klein, H.L., see Button, L.L., 271
Knapp, B., Hundt, E., Nau, U. and Küpper, H.A.
Molecular cloning, genomic structure and localization in a blood stage antigen of *Plasmodium falciparum* characterized by a serine stretch, 73
Knight, M., see Rodrigues, V., 7
Koller, B., see Huber, M., 285
Krungkrai, J., Webster, H.K. and Yuthavong, Y.
De novo and salvage biosynthesis of pteroylpentaglutamates in the human malaria parasite, *Plasmodium falciparum*, 25
Kuhns, J., see Reis, M.G., 113
Küpper, H.A., see Knapp, B., 73
Kwan-Lim, G.-E., see Maizels, R.M., 213

Lensen, T.H.W., see Delves, C.J., 105
Lillehoj, H.S., see Jenkins, M.C., 153
Lindmark, D.G., see Jarroll, E.L., 121

Maizels, R.M., Gregory, W.F., Kwan-Lim, G.-E. and Selkirk, M.E.
Filarial surface antigens: the major 29 kilodalton glycoprotein and a novel 17–200 kidodalton complex filarial surface antigens from adult *Brugia malayi* parasites, 213
Maizels, R.M., see Selkirk, M.E., 229
Mama, K., see Diffley, P., 1
Manning, P., see Jarroll, E.L., 121
Marr, J.J., see Goad, L.J., 179
Marshall, V.M., see Favaloro, J.M., 297
Matile, H., see Delves, C.J., 105
McBride, J.S., see Clark, J.T., 15
McGuire, T.C., see Barbet, A.F., 191
McMaster, W.R., see Button, L.L., 271
Meadows, H., see Rodrigues, V., 7
Medina-Acosta, E., see Button, L.L., 271
Merritt, Jr., J.W., see Aggarwal, A., 39
Mirelman, D., see Huber, M., 285
Myler, P.J., see Barbet, A.F., 191

Nakamura, Y., see Watanabe, J.-i., 163
Nash, T.E., see Aggarwal, A., 39
Nau, U., see Knapp, B., 73
Nielsen, L., see Selkirk, M.E., 229
Nonnengässer, C., see Bülow, R., 85

Ostoa-Saloma, P., Cabrera, N., Becker, I and Perez-Montfort, R.
Proteinases of *Entamoeba histolytica* associated with different subcellular fractions, 133
Overath, P., see Bülow, R., 85

Paindavoine, P., Zampetti-Bosseler, F., Coquelet, H., Pays, E. and Steinert, M.
Different allele frequenties in *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* populations, 61
Partono, F., see Selkirk, M.E., 229
Pays, E., see Paindavoine, P., 61
Perez-Montfort, R., see Ostoa-Saloma, P., 133
Perlmann, P., see Franzén, L., 201
Pettersson, U., see Franzén, L., 201
Ponnudurai, T., see Delves, C.J., 105
Prichard, R.K., see Tang, L., 145

Reis, M.G., Kuhns, J., Blanton, R. and Davis, A.H.
Localization and pattern of expression of a female specific mRNA in *Schistosoma mansoni*, 113
Revel, M., see Huber, M., 285
Ridley, R.G., see Delves, C.J., 105
Rodrigues, V., Chaudhri, M., Knight, M., Meadows, H., Chambers, A.E., Taylor, W.R., Kelly, C. and Simpson, A.J.G.
Predicted structure of a major *Schistosoma mansoni* eggshell protein, 7
Rozenblatt, S., see Huber, M., 285
Russell, D.G., see Button, L.L., 271

Sayers, G., see Selkirk, M.E., 229
Scaife, J.G., see Delves, C.J., 105
Schmatz, D.M., Baginsky, W.F. and Turner, M.J.
Evidence for and characterization of a mannitol cycle in *Eimeria tenella*, 263
Scholler, J.K., see Aline, Jr., R.F., 169
Selkirk, M.E., Nielsen, L., Kelly, C., Partono, F., Sayers, G. and Maizels, R.M.
Identification, synthesis and immunogenicity of cuticular collagens from the filarial nematodes *Brugia malayi* and *Brugia pahangi*, 229
Selkirk, M.E., see Maizels, R.M., 213
Simpson, A.J.G., see Rodrigues, V., 7
Sims, P.F.G., see Hyde, J.E., 247
Snewin, V.A., see Hyde, J.E., 247
Snounou, G., see Viriyakosol, S., 93
Steinert, M., see Paindavoine, P., 61
Stuart, K., see Aline, Jr., R.F., 169

Tanabe, K., see Watanabe, J.-i., 163
Tang, L. and Prichard, R.K.
Characterization of tubulin from *Brugia malayi* and *Brugia pahangi*, 145
Taylor, W.R., see Rodrigues, V., 7
Turner, M.J., see Schmatz, D.M., 263

Viriyakosol, S., Snounou, G. and Brown, K.N.
The use of a DNA probe for the differentiation of rodent malaria strains and species, 93

Wahlgren, M., see Franzén, L., 201
Wåhlin, B., see Franzén, L., 201
Watanabe, J.-i., Hori, H., Tanabe, K. and Nakamura, Y.
Phylogenetic association of *Pneumocystis carinii* with the 'Rhizopoda/Myxomycota/Zygomycota group' indicated by comparison of 5S ribosomal RNA sequences, 163
Webster, H.K., see Krungkrai, J., 25
Wigzell, H., see Franzén, L., 201
Williams, R.O., see Barbet, A.F., 191
Wilson, C.F., see Clark, J.T., 15
Yuthavong, U., see Krungkrai, J., 25

Zampetti-Bosseler, F., see Paindavoine, P., 61

Subject Index

Actin, 247
β-*N*-Acetylgalactosaminidase, 121
Amastigote, 179
Amino hexose, 121
Antigenic variation, 39, 191
Asparagine-rich, 201

Benzimidazole, 145
Biomphalaria glabrata, 49
Brugia, 229
Brugia malayi, 145, 213
Brugia pahangi, 145, 213

Calorimetry, 57
Carbohydrate, 121
Carbohydrate metabolism, 263
Cell cycle, 1
Chemotherapy, 179
Chromosome location, 271
Circular episome, 285
Codon match index, 247
Collagen, 229
Common proteins, 49
Cross reactivity, 229
Cuticle, 213
Cuticle structure, 229
Cysteine-rich protein, 39
Cytospectrophotometry, 1

De novo folate synthesis, 25
Diagnosis, 105
Differentiation, 85
Dihydrofolate reductase-thymidylate synthetase, 247
DNA hybridisation, 93
cDNA, 153
cDNA sequence, 297
rDNA organization, 285
25S rDNA sequence, 285

Eggshell protein, 7
Eimeria, 153
Eimeria tenella, 263
Electron microscopy, 285
Entamoeba histolytica, 133, 285
Epimastigote, 179
Exon-intron pattern, 73

Female specific cDNA, 113
Fluconazole, 179
Fluorescent probe, 57
Folate biosynthesis, 25

Galactosamine, 121
Gene evolution, 191
Gene expression, 113
Gene localization, 113
Gene organization, 271
Gene sequence, 7
Gene structure, 73
Genetic exchange, 61
Giardia, 121
Giardia lamblia, 39
Glycosylation, 213
Growth rate, 1

Hexokinase, 263
Hydrophobic chromatography, 133

In situ hybridization, 113
Itraconazole, 179

Ketoconazole, 179

Leishmania, 271

Major surface glycoprotein, 271
Malaria, 93, 105, 201
Mannitol, 263
Mannitol cycle, 263
Mannitol dehydrogenase, 263
Mannitol-1-phosphatase, 263
Mannitol-1-phosphate dehydrogenase, 263
Membrane, 57
Metabolic labeling, 39
Monoclonal antibody, 15, 201

Nematode, 145, 213
Nucleic acid hybridisation, 247

Octapeptide-repeat antigen, 297
Oligonucleotide probe, 247

Peptide map, 213
Plasmodium berghei, 93
Plasmodium chabaudi, 93
Plasmodium falciparum, 15, 25, 73, 105, 201, 297
Plasmodium vinckei, 93
Plasmodium yoelii, 93
Pneumocystis carinii, 163
Polyols, 263
Polyribosome, 169
Predicted structure, 7
Proteinase, 133
Protozoan, 39
Protozoan cyst wall, 121
Pteroylpentaglutamate, 25

Qinghaosu, 57

Recombinant antigen, 153
Reducing agent, 213
Repetitive DNA, 105, 201

Restriction fragment length polymorphism, 61
5S Ribosomal RNA, 163
Rodent, 93

Salvage pathway, 25
Schistosoma mansoni, 7, 49
Serine protein, 73
Sporozoite, 153
Sterol, 179
Subcellular fraction, 133
Substrate gel electrophoresis, 133
Surface coat, 85
Surface glycoprotein, 15, 191
Surface iodination, 39
Synthesis, 229

Taxonomy, 93, 163
Transcription, 169, 271
Triton X-114 phase separation, 133
Trypanosoma b. gambiense speciation, 61
Trypanosoma brucei, 85, 169, 191
Trypanosoma brucei gambiense, 1
Trypanosoma cruzi, 179
Trypanosome, 61
Trypomastigote, 179
Tubulin, 247
Tubulin isoform, 145
Turnover, 85

Variant surface glycoprotein, 85, 169
Vitelline cell, 113